
# Nichtinvasive Ablation von Kammertachykardien



Arash Arya





### Disclosures:

**NONE** 

### Herbsttagung Schkopau 15 - 16 November 2019



Home

#### Non Invasive Ablation of Ventricular Tachycardias

Current perspective and future direction Nichtinvasive Ablation von Kammertachyka[...] PDF-Dokument [24.8 MB]

#### CardioSummit Berlin 04 - 05 October 2019



#### **Can VT Ablation Prevent Sudden Cardiac Death?**

**Current Perspective and Future Direction** Can VT Ablation Prevent SCD.pdf PDF-Dokument [40.6 MB]

### Patienten-Semonar: 25 Jahren Herzzentrum Leipzig, 28.09.2019



#### Bringen Medikamente die Libido durcheinander?

Nur für Patienten geeignet Sexuelle Dysfunktion bei Herzpatienten.p[...] PDF-Dokument [9.3 MB]

### 2019 HRS/EHRA Focused Update on Optimal ICD Programming



#### **Optimal ICD Programming and Testing**

2019 HRS/EHRA Update optimal ICD programming and Testing.pdf PDF-Dokument [1.7 MB]

### Überblick:

### Nichtinvasive Ablation von Kammertachykardien



**Aktuelle Studien** 





**Meine Vision** 


# VT refractory to conventional Ablation:


- Increase Power
- Change Catheter
- Using Half-Saline Irrigation
- Bipolar Ablation
- Alcohol Ablation
- Bilateral surgical sympathectomy
- Stereotactic Radio-Surgery
- Electroporation

### **Supplemental Figure S2:** Thresholds of Various Cell Types to PFA









Comput Biol Med. 2018 Nov 1;102:376-380

 Table 1
 Shown are key differences between conventional catheter ablation and stereotactic arrhythmia radioablation


	Catheter ablation	Stereotactic arrhythmia radioablation
Energy source	Radiofrequency energy, cryoablation, others	Ionizing radiation
Pathophysiology of tissue destruction	Tissue heating or freezing leading to necrosis	Likely apoptosis and micro-vascular injury
Time course for lesion formation	Seconds to minutes	Days to months
Tissues that can be targeted	Local injury to tissue in direct contact or close-by proximity to ablative catheter; therefore, tissues accessible by catheter can be ablated	Any tissue is theoretically ablatable
Lesion geometry	Point lesions or planar lesions in contact with catheters	3-dimensional volume
Benefits	<ul><li>Established technology</li><li>Near-immediate effect</li></ul>	<ul> <li>Noninvasive</li> <li>Any tissue is potentially ablatable, including deep targets</li> <li>3D ablation volume can be created</li> </ul>
Risks	<ul><li>Invasive risks</li><li>Collateral injury</li></ul>	<ul><li>Radiation injury</li><li>Unknown late side effects</li></ul>

Curr Cardiol Rep (2017) 19:79

### ORIGINAL ARTICLE

## Noninvasive Cardiac Radiation for Ablation of Ventricular Tachycardia

Phillip S. Cuculich, M.D., Matthew R. Schill, M.D., Rojano Kashani, Ph.D., Sasa Mutic, Ph.D., Adam Lang, M.D., Daniel Cooper, M.D., Mitchell Faddis, M.D., Ph.D., Marye Gleva, M.D., Amit Noheria, M.B., B.S., Timothy W. Smith, M.D., D.Phil., Dennis Hallahan, M.D., Yoram Rudy, Ph.D., and Clifford G. Robinson, M.D.




From April through November 2015, five patients with high-risk, refractory VT underwent treatment.

The mean noninvasive ablation time was 14 minutes (range, 11 to 18).

During the 3 months before treatment, the patients had a combined history of 6577 episodes of ventricular tachycardia.

After the 6-week blanking period, there were 4 episodes of ventricular tachycardia over the next 46 patient-months, for a reduction from baseline of 99.9%.







### **CLINICAL RESEARCH**

Ablation for ventricular tachycardia

## Stereotactic radiosurgery for ablation of ventricular tachycardia

Radek Neuwirth<sup>1</sup>, Jakub Cvek (1) <sup>2\*</sup>, Lukas Knybel<sup>2</sup>, Otakar Jiravsky<sup>3</sup>, Lukas Molenda<sup>2</sup>, Michal Kodaj<sup>3</sup>, Martin Fiala<sup>1</sup>, Petr Peichl<sup>4</sup>, David Feltl<sup>5</sup>, Jaroslav Januška<sup>3</sup>, Jan Hecko<sup>3</sup>, and Josef Kautzner<sup>4</sup>








**Figure I** Reduction of sustained VT before and after treatment. (A) Significant reduction of sustained VT after radiosurgery is shown. (B) The total number of episodes of sustained VT in each of the 10 study patients, for three consecutive months before treatment (-1.Q) and continuing for 3 months periods after treatment (1.Q-18.Q) is shown. VT, ventricular tachycardia.


### What's new?


- This is case series with the longest follow-up ever on safety and efficacy of stereotactic body radiotherapy in patients with structural heart disease and recurrent ventricular tachycardia (VT).
- Radiotherapy in our patients resulted in a significant decrease of sustained VTs.
- Two our patients showed practically no treatment effect, and in other two patients the effect of radiotherapy was significantly delayed.
- The only acute radiation-related toxicity observed in our study was nausea.
- One possible late cardiac radiation-related toxicity Grade 3 was found 17 months after radiosurgery.
















Die Zukunft der VT-Ablation



künstliche Intelligenz

### Screening for cardiac contractile dysfunction using an artificial intelligence-enabled electrocardiogram


Zachi I. Attia¹, Suraj Kapa¹, Francisco Lopez-Jimenez¹, Paul M. McKie 📵¹, Dorothy J. Ladewig², Gaurav Satam², Patricia A. Pellikka 📵¹, Maurice Enriquez-Sarano¹, Peter A. Noseworthy 📵¹, Thomas M. Munger¹, Samuel J. Asirvatham¹, Christopher G. Scott³, Rickey E. Carter 📵⁴ and Paul A. Friedman 🔞¹\*

Nature Medicine I VOL 25 I 70 JANUARY 2019 I 70-74 I www.nature.com/naturemedicine

Using paired 12-lead ECG and echocardiogram data, including the LVEF, from 44,959 patients at the Mayo Clinic, we trained a convolutional neural network to identify patients with ventricular dysfunction, defined as ejection fraction ≤ 35%, using the ECG data alone.

When tested on an independent set of 52,870 patients, the network model yielded values for the area under the curve, sensitivity, specificity, and accuracy of 0.93, 86.3%, 85.7%, and 85.7%, respectively.

In patients without ventricular dysfunction, those with a positive Al screen were at 4 times the risk (hazard ratio, 4.1; 95% confidence interval, 3.3 to 5.0) of developing future ventricular dysfunction compared with those with a negative screen.



An artificial intelligence-enabled ECG algorithm for the identification of patients with atrial fibrillation during sinus rhythm: a retrospective analysis of outcome prediction

Zachi I Attia\*, Peter A Noseworthy\*, Francisco Lopez-Jimenez, Samuel J Asirvatham, Abhishek J Deshmukh, Bernard J Gersh, Rickey E Carter, Xiaoxi Yao, Alejandro A Rabinstein, Brad J Erickson, Suraj Kapa, Paul A Friedman

LANCET: http://dx.doi.org/10.1016/S0140-6736(19)31721-0

... Including all ECGs acquired during the first month of each patient's window of interest (i.e., the study start date or 31 days before the first recorded atrial fibrillation ECG) increased the AUC to 0.90 (0.90–0.91), sensitivity to 82.3% (80.9–83.6), specificity to 83.4% (83.0–83.8), F1 score to 45.4% (44.2–46.5), and overall accuracy to 83.3% (83.0–83.7).





Figure 3: ROC curves for the convolutional neural networks on the testing dataset



**Fig. 3 | The virtual medical coach model with multi-modal data inputs and algorithms to provide individualized guidance.** A virtual medical coach that uses comprehensive input from an individual that is deep learned to provide recommendations for preserving the person's health. Credit: Debbie Maizels/Springer Nature


*Nature Medicine* **volume 25**, pages44–56 (2019)

Rapid 12-lead automated localization method: Comparison to electrocardiographic imaging (ECGI) in patient-specific geometry

Shijie Zhou, PhD <sup>a,b</sup>, B. Milan Horáček, PhD <sup>b</sup>, James W. Warren, BSc <sup>d</sup>, Amir AbdelWahab, MD <sup>a,c</sup>, John L. Sapp, MD <sup>a,c</sup>,\*


Journal of Electrocardiology. 2019: accepted manuscript


With 10 training pacing sites ... demonstrated that the automated method achieved localization error of <5mm for the VT-exit site



### Fast personalized electrophysiological models from computed tomography images for ventricular tachycardia ablation planning

The authors aim at building such a pipeline from computed tomography (CT) images to personalized cardiac electrophysiology (EP) model.








Nicolas Cedilnik et al. Europace (2019) In Press.


















Der beste Weg, die Zukunft vorherzusagen, ist, sie zu gestalten.

Peter Drucker